

A Review of Biogenic Amines and Polyamines in Beer

Pavel Kaláč^{1,2} and Martin Křížek¹

ABSTRACT

J. Inst. Brew. 109(2), 123–128, 2003

The presence of biogenic amines and polyamines in foods and alcoholic beverages is important from both toxicological and technological points of view. High amounts of these compounds can lead to health problems and accurate methods of determination, as well as knowledge on where the compounds originate and how they can be controlled at the lowest levels, are important in the food and beverage industry. In brewing, the types of amines are dependent on the raw materials in the beverage, as well as the method of brewing, and any microbial contamination that may have occurred during the brewing process or during storage. Studies looking at biogenic amine and polyamine levels in various beers have been carried out by a number of researchers in Europe, Brazil, Canada and Cuba. This paper reviews the work from studies carried out previously and summarizes the values found by the various research groups. Methods of analysis for the amines including HPLC, HPTLC and enzyme immunoassays are reviewed.

Key words: Beer, biogenic amines, lactic acid bacteria, polyamines, tyramine.

INTRODUCTION

Biogenic amines (BAs) form a group of undesirable natural components widespread in foods and beverages, e.g. scombroid fish, meat and meat products, cheeses, vegetable products and wine (for reviews see^{39,41}). Beer has also been reported as a possible health risk for some consumers due to BAs intake⁹. These high biogenic amine intake levels are not usually caused by a high amine content in individual beers, but rather by a very high beer consumption during a very short time interval. Although annual statistical beer consumption exceeds 100 L/capita/year in several countries, the consumption referred to here is several times higher in these particular individuals. Beer is now considered as a source of the dietary polyamines, putrescine, spermidine and spermine. These regulate cellular proliferation and differentiation^{2,33}.

BAs in foods and beverages arise mainly from the bacterial decarboxylation of the corresponding amino acids

(Fig. 1). Thus, histamine (HI), tyramine (TY), tryptamine (TR), 2-phenylethylamine (PEA), agmatine (AGM) and cadaverine (CAD) are formed from histidine, tyrosine, tryptophan, phenylalanine, arginine and lysine, respectively. Putrescine (PUT) can arise both from ornithine and agmatine. Spermidine (SPD) and spermine (SPM) are formed biochemically from putrescine by the attachment of aminopropyl moieties catalysed by spermidine synthase and spermine synthase⁴⁵.

Excessive intake of dietary BAs can cause several deleterious effects. Psychoactive amines can affect the neural transmitters in the central nervous system. Vasoactive amines can act directly or indirectly on the vascular system as vasoconstrictors (namely TY) or vasodilators (HI).




Fig. 1. Structures of some common biogenic amines.

¹ University of South Bohemia, Faculty of Agriculture, Department of Chemistry, 370 05 České Budějovice, Czech Republic
² Corresponding author. E-mail: kalac@zf.jcu.cz

TABLE I. Histamine content (mg/L) in beer.

Year	Country	Number of samples	Mean	SD	Range	References
1989	Spain	17	0.7	—	0.3–1.8	22
1989	7 European countries	48	1.4	—	0.3–17.0	22
	Spain	35 (common)	0.8	1.8	0.35–1.8	23
		20 (special)	1.2	1.4	0.3–4.9	
1991	Spain	30	0.7	0.4	—	14
1996	16 European countries	195	1.2	2.4	<0.3–21.6	18
1997	Czech Rep.	78	0.6	1.1	<0.3–4.7	24
1999	Brazil	91	0.2	0.3	<0.2–1.5	11

SD = standard deviation

TABLE II. Tyramine content (mg/L) in beer.

Year	Country	Number of samples	Mean	SD	Range	References
1989	Cuba	104	3.5–8.3 ^a	—	1.2–9.3	47
1989	Spain	17	5.8	—	1.9–24.7	22
1989	7 European countries	48	6.1	—	1.5–46.8	22
1991	Spain	35 (common)	4.7	5.2	1.6–30.6	23
		20 (special)	7.7	7.7	1.9–24.8	
1995	Italy	16	4.3	—	0.6–17.2	3
1996	Spain	30	6.8	7.4	—	14
1996	16 European countries	195	6.5	9.0	0.5–67.5	18
1997	Czech Rep.	78	6.9	5.2	<0.3–22.5	24
1999	Brazil	91	2.2	4.8	0.3–36.8	11

^a Mean values increased with increasing alcohol level.

A scale of symptoms can occur following the excessive oral intake of BAs. The particular risk strongly depends on detoxification efficiency, which can vary considerably between individuals and is affected by several factors. Normal intakes of BAs are metabolised in the intestinal tract by a fairly efficient detoxification system based on the activities of monoamine oxidase (MAO; EC 1.4.3.4) and diamine oxidase (DAO; EC 1.4.3.6).

The limited literature data on BAs content and formation in beer was reviewed by Stratton et al. in 1991⁴³. The aim of this article is to review the information published since that review.

HEALTH RISKS OF BIOGENIC AMINES FOR BEER CONSUMERS

Hypertensive crises have been observed in patients treated with drugs inhibiting monoamine oxidase (used mainly in psychiatry) after beer consumption^{31,35,40,44}. The adverse effects were observed from both draft and non-alcoholic beers and were attributed to TY. TY intake as low as 6 mg within a 4-hour period or beers with TY content over 10 mg/L were considered as dangerous for these particular patients⁴⁴. Alcohol and probably some other BAs present in beer can potentiate TY effects. However, no risk was reported for healthy consumers.

Beer has been reported as a trigger for headaches with patients susceptible to migraines³⁶. Histamine in alcoholic drinks was capable of triggering of allergic and allergic-like adverse responses, however, wine has been a more common source than beer⁴⁶. Secondary amines (AGM, SPD and SPM) can react with nitrites to form nitrosamines, but no literature was found dealing with this particular reaction in beer.

BIOGENIC AMINES – CONTENT IN BEERS

Data on BAs in beers, analysed over the past few years has been collated chronologically in Tables I–IV. The results from surveys in several European countries, Brazil and Cuba are given for beers of different types. Although different analytical methods were used, the reported levels are comparable.

The nine determined amines can be divided into two groups. Group one includes PUT, SPD, SPM and AGM and can be considered as natural beer constituents primarily originating from the malt, whilst Group two, mainly HI, TY and CAD, usually indicate the activities of contaminating lactic acid bacteria during brewing¹⁸.

Agmatine (Table IV), a compound not routinely determined probably due to its minimal adverse effects for man, was the prevailing amine in the beers tested, with mean values around 10 mg/L^{11,14,18}. TY and PUT (Tables II and III) were amines at relatively high levels, while the levels of the others, including HI, have commonly been lower and are probably of lower toxicological importance. Thus, TY has been the main biogenic amine identified in beer that causes the adverse effects described earlier and high levels were reported by Tailor et al.⁴⁴.

Similar levels of BAs were found in both alcoholic and non-alcoholic beers^{3,11,18,24}, suggesting that the production methods for non-alcoholic beers do not remove amines. No significant correlations were found between TY or HI levels and alcohol content, or extract of original wort^{22,24}. However, higher levels of the both amines were observed in beers with a high total acidity²², which indicated activity of lactic acid bacteria.

Spontaneously fermented Belgian beers and top-fermented beers showed the highest levels of TY and HI and

TABLE III. Polyamine (putrescine, spermidine and spermine) content (mg/L) in beer.

	Year	Country	Number of samples	Mean	SD	Range	References
Putrescine	1992	Belgium	11	4.6	—	2.6–8.4	8
	1995	Italy	16	2.9	—	1.4–4.2	3
	1996	Spain	30	4.5	1.7	—	14
	1996	16 European countries	195	4.8	2.3	1.5–15.2	18
	1997	Czech Rep.	78	8.8	7.1	<0.3–30.7	24
	1999	Brazil	91	3.9	1.4	0.9–9.8	11
	2002	Poland	27	1.8	0.8	0.6–3.6	42
Spermidine	1992	Belgium	11	0.3	—	<0.2–0.8	8
	1995	Italy	16	0.7	—	0.3–1.4	3
	1996	Spain	30	0.8	1.0	—	14
	1996	16 European countries	195	0.7	1.0	<0.3–6.8	18
	1999	Brazil	91	0.7	0.8	<0.2–6.0	11
Spermine	1995	Italy	16	ND	—	<0.2	3
	1996	Spain	30	0.2	0.4	—	14
	1996	16 European countries	195	0.3	0.7	<0.3–3.9	18
	1996	Brazil	91	0.3	0.5	<0.2–2.1	11
	2002	Poland	27	8.4	3.6	1.2–15.2	42

TABLE IV. Cadaverine, tryptamine, 2-phenylethylamine and agmatine content (mg/L) in beer.

	Year	Country	Number of samples	Mean	SD	Range	References
Cadaverine	1992	Belgium	11	3.2	—	<0.15–13.3	8
	1995	Italy	16	—	—	0.2–0.8	3
	1996	Spain	30	0.7	0.4	—	14
	1996	16 European countries	195	2.4	6.1	<0.3–39.9	18
	1997	Czech Rep.	78	12.9	12.3	<0.3–49.1	24
	1999	Brazil	91	0.5	0.4	<0.2–2.6	11
	Tryptamine	1995	Italy	16	<1.0	—	ND–2.6
1996		Spain	26	1.6	2.1	—	14
1996		16 European countries	195	0.4	1.0	<0.3–5.4	18
1997		Czech Rep.	78	1.2	1.5	<0.3–9.7	24
1999		Brazil	91	0.5	1.5	<0.35–10.1	11
2-Phenylethyl-amine	1995	Italy	16	1.0	—	0.5–1.6	3
	1996	Spain	30	0.3	0.1	—	14
	1996	16 European countries	195	0.4	0.8	<0.3–8.3	18
	1999	Brazil	91	0.1	0.2	<0.2–1.7	11
Agmatine	1996	Spain	30	9.3	4.5	—	14
	1996	16 European countries	195	10.5	5.8	0.5–40.9	18
	1999	Brazil	91	10.9	7.0	2.1–46.8	11
	1999	Belgium	150	—	—	4.0–30	32

also of PEA and TR^{10,32}. These types of beers represent the highest toxicological risk for patients treated with MAO inhibitors. Lorent et al.³² have proposed a safety limit of 20 mg/L for the sum of the TY + HI + CAD + PEA contents.

Low levels of TY, HI, PUT, CAD, PEA and TR, but not exceeding 0.5 mg/L of the individual amines, were observed in a survey of local Nigerian ginger beer and beers produced from sorghum, maize and plantain²⁹.

The question of whether increased BAs levels affect the sensorial quality of beer is still unresolved.

BIOGENIC AMINES IN RAW MATERIALS

Data on BAs levels in malt, hops or hop derivatives (pellets, extract) as well as pitching yeast is very limited^{13,19,24}. No amines should be present in production water. PUT, SPD, SPM and AGM are present in malt and yeast at higher levels than in hops. The levels of the other amines are also low in these raw materials. Taking into account the low levels of hops and yeast in beer, it can be

said that malt is the main source of the four aforementioned amines in beer. Please note that a portion of these amines is removed in the spent grain²⁴.

HI, PEA, TR and CAD levels were seen to increase slowly during a five-day barley germination, and a increase of 3–5.5 mg/kg/day was observed for PUT, SPD, SPM and AGM. The TY increase was lower¹⁹. Malting conditions such as germinating intensity, kiln temperature and barley variety will all affect the final amine levels in malt¹³. Amines were not detected in the rice used as an adjunct cereal¹⁹.

BIOGENIC AMINES – FORMATION DURING BREWING

The greatest TY^{6,12,19,23,24}, HI^{12,24} and CAD²⁴ formation was observed during the main fermentation. Levels of PUT^{13,24} and AGM¹³ decreased starting from the mashing process. The bottom yeast *Saccharomyces cerevisiae* var. *uvarum* did not appear to produce TY or HI during fermentation and moreover, yeast recycling for several fer-

mentations did not influence amine levels¹⁶. Also wild yeasts were not found to produce TY, while a significant positive relationship was observed between amine formation and lactic acid bacteria¹⁷.

Bacteria isolated during beer fermentation and distinguished by TY and TR formation were identified as *Pediococcus* spp., mainly *P. damnosus*. No *Lactobacillus* spp. were isolated. TY formation was negligible at *Pediococci* spp. counts below 4×10^3 colony-forming units (CFU) per mL, while at over 1×10^5 CFU/mL, TY levels ranged between 15–25 mg/L¹⁷. Thus, TY content has been proposed as a reliable indicator of the level of *Pediococcus* spp. contamination during a beer fermentation¹⁵. Washing the pitching yeast with phosphoric acid was an effective way to reduce the number of *Pediococci* and consequently reduce the TY content of the beer¹⁷. However, species of the genus *Lactobacillus* also participate in BAs formation. *Lactobacillus frigidus*, *L. brevis* and *L. brevis* were reported as amine-forming beer contaminants⁷.

Different strains of lactic acid bacteria found in foods have been reported to differ widely in their ability to produce BAs⁴. This situation could also exist in brewery lactic acid populations and may explain the differences existing in TY, and also HI contents, among breweries and among batches within one type of beer from the same brewery.

A direct relationship was not observed between the free tyrosine levels in wort and TY formation during fermentation. Content of this amino acid does not seem to be a critical factor in TY formation²⁰.

Amines can also be formed by the activity of lactic acid bacteria during the storage of beer in bottles, cans or kegs prior to consumption. Secondary fermentation in some bottles of special beers resulted in elevated BAs levels³². However, most of the beers studied were pasteurised. A significant increase in TY and HI contents that was observed in one pasteurised beer stored for several weeks²⁴, was probably due to under-pasteurisation. Considerable increases of TY, and to a lesser extent HI, were found in adequately pasteurised beers that were inoculated post pasteurisation with mixed *Lactobacillus* spp. or *Pediococcus* spp. isolated from spoiled beer, and stored at 28°C until haze formation commenced. *Lactobacilli* were reported to be more effective amine producers than *Pediococci*. Minimum changes in PUT, CAD and SPD levels were observed²⁵.

Lactobacillus paracasei isolated from a bottled wheat beer produced no HI, PUT or CAD during cultivation in malt medium at 30°C for 7 days, or in beer in opened bottles. In contrast, *Lactobacillus buchneri* cultivated as a negative control, increased the HI content in beer from 15 to 65 mg/L, while the PUT and CAD content did not change⁴⁹. The HI content in beer has been proposed as a good indicator of brewing hygiene since it does not originate from the malt¹³.

ANALYTICAL METHODS

Methods for the measurement of biogenic amines in foods are not numerous and only a few of these methods have been applied to beer. The first procedures developed

focused on the most toxicologically important amines – HI and TY. Izquierdo-Pulido et al.²³ used two extraction methods. HI was extracted to n-butanol after sample alkalisation, re-extracted with hydrochloric acid and determined spectrofluorimetrically as a fluorescent complex with *o*-phthalaldehyde (OPA). Extraction of TY similarly originated in an alkaline sample and was carried out with ethyl acetate. After re-extraction with hydrochloric acid, the tyramine complex with α -nitroso- β -naphthol was measured using spectrofluorimetry. As other biogenic amines are also of interest, the same authors²¹ developed a comprehensive and more selective method. Ten amines – HI, TY, serotonin, PEA, TR, CAD, PUT, AGM, SPD and SPM – were separated in one run using ion-pair chromatographic partition on a reversed-phase column. After separation, the amines were detected as fluorescent derivatives with *o*-phthalaldehyde. Determination limits of the method ranged from 0.3 to 0.4 mg/L, except for serotonin and SPM, which were slightly higher.

Though post-column derivatization is more common when using *o*-phthalaldehyde, Petridis et al.³⁷ described a simple and selective HPLC method for beers and other beverages with pre-column OPA derivatization. Recoveries for HI, TY, PUT, CAD, TR and PEA were higher than 95%. The detection limits of the BAs ranged between 5–25 μ mol/L.

In recent years, Arlorio et al.¹ have combined the extraction pre-separation step with ion-pair chromatography. The extraction steps were carried out by either butanol or an ionic exchanger. The authors concluded that many interfering compounds were eliminated by the pre-separation. However, this procedure appears to be laborious and detection limits were not given.

Alternatively to the detection of post-column derivatives, another approach was the separation of derivatives prepared prior to high performance liquid chromatography. Buiatti et al.³ separated eight amines (HI, TY, PEA, TR, CAD, PUT, SPD and SPM). Amine derivatives with dansyl chloride were eluted from a reversed-phase column by an acetonitrile/water solution at pH 7. The gradient of acetonitrile concentration in the mobile phase was 65–90% during 0–6 min. For detection, the UV detector was set at 254 nm.

Derivatives with dansyl chloride were also separated by a high performance thin-layer chromatographic method (HPTLC). Microwave assisted dansylation reduced analysis time and the detection limits of the amines ranged between 0.28–0.39 ng³⁰.

Another agent was used for pre-column derivatization of beer amines by Křížek and Hlavatá²⁷. Benzoyl chloride forms stable derivatives with biogenic amines and these derivatives were extracted from beer samples adjusted to pH 6.0 by a phosphate buffer solution. The determination limit was 0.3 mg/L. More recently, the reagent *p*-nitrobenzoyloxycarbonyl chloride (PNZ-Cl) was used for pre-column derivatization of biogenic amines in fermented beverages and vinegars²⁶. This method was applied to the determination of PEA, TR, PUT, CAD, HI, TY, SPD, SPM and serotonin. Recoveries of BAs ranged from 78% to 93%.

Another technique for beer amine determination is the amino acid analyser. This procedure employs a potassium

citrate buffer system and a post-column ninhydrin reaction¹³.

HI in beers has also been determined by capillary zone electrophoresis with amperometric detection⁴⁸. This method did not require any sample clean-up procedures, but used a detector not ranked among standard detector types.

Very promising techniques involve enzyme immunoassay methods or selective enzyme biosensors. For the determination of HI in beers a competitive enzyme immunoanalysis has been developed⁵. This method is very selective and not affected by either HI, TY or their various derivatives. The proof limit of this method is 7 ng/mL. A more versatile enzyme sensor for the simultaneous determination of three BAs (HI, TY and PUT) was designed by Lange and Wittmann²⁸. However, the lower detection limits with this sensor (range between 5–10 mg/kg) appeared to lack sensitivity.

CONCLUSIONS

There is currently much work ongoing globally as researchers try to better understand the impact of biogenic amines and polyamines in our foods^{34,38}. It is clear that in beer, bacterial contamination is a key source of these compounds and that this is a source that can be controlled to ensure that the beer meets all quality and safety standards. In addition better methods of analysis for these compounds are being developed which allows for better monitoring.

ACKNOWLEDGEMENTS

This review was prepared within research project 525/02/1077 financed by the Grant Agency of the Czech Republic and within the framework of the COST 922 project.

REFERENCES

1. Arlorio, M., Coisson, J.D. and Martelli, A., Extraction methods for biogenic amines in wine and beer. *Ital. J. Food Sci.*, 1999, **11**, 355–360.
2. Bardócz, S., Duguid, T.J., Brown, D.S., Grant, G., Pusztai, A., White, A. and Ralph, A., The importance of dietary polyamines in cell regeneration and growth. *Br. J. Nutr.*, 1995, **73**, 819–828.
3. Buiatti, S., Boschelle, O., Mozzon, M. and Battistutta, F., Determination of biogenic amines in alcoholic and non-alcoholic beers by HPLC. *Food Chem.*, 1995, **52**, 199–202.
4. Bover-Cid, S. and Holzapfel, W.H., Improved screening procedure for biogenic amine production by lactic acid bacteria. *Int. J. Food Microbiol.*, 1999, **53**, 33–41.
5. Čepička, J., Rychetský, P., Hochel, I., Strejček, F. and Rauch, P., Immunoenzymic determination of histamine in beer. *Monatsschr. Brauwiss.*, 1992, **42**, 161–164 (in German).
6. Cerutti, G., Finoli, C. and Vecchio, A., Non-volatile amine biogenesis from wort to beer. *Monatsschr. Brauwiss.*, 1989, **42**, 246–248.
7. Donhauser, S., Wagner, D. and Geiger, E., Biogenic amines. Importance, intake and evaluation. *Brauwelt*, 1992, **27**, 1272–1280 (in German).
8. Dumont, E., De Geeter, H. and Huyghbaert, A., Presence and formation of biogenic amines in local Belgian beers. *Med. Fac. Landbouww. Univ. Gent*, 1992, **57/4b**, 1911–1913.
9. Gardner, D.M., Shulman, K.I., Walker, S.E. and Tailor, S.A.N., The making of a user friendly MAOI diet. *J. Clin. Psychiat.*, 1996, **57**, 99–104.
10. Gasarasi, G., Kelgtermans, M., Verstrepen, K.J., Van Roy, J., Delvaux, F.R. and Derdelinckx, G., Occurrence of biogenic

amines in beer: causes and proposals of remedies. *Monatsschr. Brauwiss.*, 2003, **56**, (3–4), 58–63.

11. Glória, M.B.A. and Izquierdo-Pulido, M., Levels and significance of biogenic amines in Brazilian beers. *J. Food Compos. Anal.*, 1999, **12**, 129–136.
12. Gorinstein, S., Zenser, M., Vargas-Albores, F., Ochoa, J.-L., Paredes-Lopez, O., Scheler, Ch., Salnikow, J., Martin-Belloso, O. and Trakhtenberg, S., Proteins and amino acids in beers, their content and relationships with other analytical data. *Food Chem.*, 1999, **67**, 71–78.
13. Halász, A., Baráth, Á. and Holzapfel, W.H., The biogenic amine content of beer; the effect of barley, malting and brewing on amine concentration. *Z. Lebensm. Unters. Forsch. A*, 1999, **208**, 418–423.
14. Izquierdo-Pulido, M., Albalá-Hurtado, S., Mariné-Font, A. and Vidal-Carou, M.C., Biogenic amines in Spanish beers: differences among breweries. *Z. Lebensm. Unters. Forsch.*, 1996, **203**, 507–511.
15. Izquierdo-Pulido, M., Carceller-Rosa, J.-P., Mariné-Font, A. and Vidal-Carou, M.C., Tyramine formation by *Pediococcus* spp. during beer fermentation. *J. Food Protect.*, 1997, **60**, 831–836.
16. Izquierdo-Pulido, M., Font-Fábregas, J. and Vidal-Carou, C., Influence of *Saccharomyces cerevisiae* var. *uvarum* on histamine and tyramine formation during beer fermentation. *Food Chem.*, 1995, **54**, 51–54.
17. Izquierdo-Pulido, M., Font-Fabregas, J., Carceller-Rosa, J.-P., Mariné-Font, A. and Vidal-Carou, C., Biogenic amine changes related to lactic acid bacteria during brewing. *J. Food Protect.*, 1996, **59**, 175–180.
18. Izquierdo-Pulido, M., Hernández-Jover, T., Mariné-Font, A. and Vidal-Carou, M.C., Biogenic amines in European beers. *J. Agric. Food Chem.*, 1996, **44**, 3159–3163.
19. Izquierdo-Pulido, M., Mariné-Font, A. and Vidal-Carou, M.C., Biogenic amines formation during malting and brewing. *J. Food Sci.*, 1994, **59**, 1104–1107.
20. Izquierdo-Pulido, M., Mariné-Font, A. and Vidal-Carou, M.C., Effect of tyrosine on tyramine formation during beer fermentation. *Food Chem.*, 2000, **70**, 329–332.
21. Izquierdo-Pulido, M., Vidal-Carou, M.C. and Mariné-Font, A., Determination of biogenic amines in beers and their raw materials by ion-pair liquid chromatography with postcolumn derivatization. *J. AOAC Int.*, 1993, **76**, 1027–1032.
22. Izquierdo-Pulido, M.L., Vidal-Carou, M.C. and Mariné-Font, A., Histamine and tyramine in beers: contents and relationship with other analytical data. *J. Food Compos. Anal.*, 1989, **2**, 219–227.
23. Izquierdo-Pulido, M.L., Vidal-Carou, M.C. and Mariné-Font, A., Histamine and tyramine in beers. Changes during brewing of a Spanish beer. *Food Chem.*, 1991, **42**, 231–237.
24. Kalač, P., Hlavatá, V. and Křížek, M., Concentrations of five biogenic amines in Czech beers and factors affecting their formation. *Food Chem.*, 1997, **58**, 209–214.
25. Kalač, P., Šavel, J., Křížek, M., Pelikánová, T. and Prokopová, M., Biogenic amine formation in bottled beer. *Food Chem.*, 2002, **79**, 431–434.
26. Kirschbaum, J., Meier, A. and Bruckner, H., Determination of biogenic amines in fermented beverages and vinegars by pre-column derivatization with para-nitrobenzylloxycarbonyl chloride (PNZ-Cl) and reversed-phase LC. *Chromatographia*, 1999, **49**, 117–124.
27. Křížek, M. and Hlavatá, V., The determination of selected biogenic amines in beer. *Kvasný Prům.*, 1995, **41**, 265–269 (in Czech).
28. Lange, J. and Wittmann, C., Enzyme sensor array for the determination of biogenic amines in food samples. *Anal. and Bioanal. Chem.*, 2002, **372**, 276–283.
29. Lasekan, O.O. and Lasekan, W.O., Biogenic amines in traditional alcoholic beverages produced in Nigeria. *Food Chem.*, 2000, **69**, 267–271.
30. Linares, R.M., Ayala, J.H., Afonso, A.M. and Gonzales, V., Determination of polyamines in beers by HPTLC with fluorescence detection. *JPC–J. Planar. Chromat.*, 2001, **14**, 4–7.

31. Lippman, S.B. and Nash, K., Monoamine-oxidase inhibitor update. Potential adverse food and drug-interactions. *Drug Safety*, 1990, **5**, 195–204.
32. Loret, S., Bianchi, D., Peulen, O., Deloyer, P. and Dandriofosse, G., Biogenic amines in different types of Belgian beers: a first survey. In: COST 917: Biogenically Active Amines in Food, S. Bardocz, J. Koninkx, M. Grillo and A. White, Eds. Office for Offic. Publ. of the Eur. Commun., Luxembourg, 1999, Vol. III, pp. 36–40.
33. Milovic, V., Polyamines in the gut lumen: bioavailability and biodistribution. *Eur. J. Gastroenterol. Hepatol.*, 2001, **13**, 1021–1025.
34. Morgan, D.L.M., Hirvi, T., Dandriofosse, G., Deloyer, P. and White, A., Eds., COST 917: Biogenically active amines in food. Office for Official Publ. of the Eur. Commun., Luxembourg, 2002, Vol. VI, 205 pp.
35. Murray, J.A., Walker, J.F. and Doyle, J.S., Tyramine in alcohol-free beer. *Lancet*, 1988, **1** (8595), 1167–1168.
36. Peatfield, R.C., Relationships between food, wine, and beer-precipitated migrainous headaches. *Headache*, 1995, **35**, 355–357.
37. Petridis, K.D. and Steinhart, H., Automated precolumn derivatization with o-phthalaldehyde (OPA) – a new RP-HPLC method for the determination of biogenic amines in foods. *Z. Lebensm.-Unters. Forsch.*, 1995, **201**, 256–260 (in German).
38. Roig, A.X., Ed., Health Implications of Dietary Amines. Book of abstracts, COST 922 Project, Univ. Autonom. Barcelona (Spain), May 2003, 50 pp.
39. Shalaby, A.R., Significance of biogenic amines to food safety and human health. *Food Res. Int.*, 1996, **29**, 675–690.
40. Shulman, K., Tailor, S.A.N., Walker, S.E. and Gardner, D.M., Tap (draft) beer and monoamine oxidase inhibitor dietary restrictions. *Can. J. Psychiat.*, 1997, **42**, 310–312.
41. Silla Santos, M.H., Biogenic amines: their importance in foods. *Int. J. Food Microbiol.*, 1996, **29**, 213–231.
42. Slomkowska, A. and Ambroziak, W., Biogenic amine profile of the most popular Polish beers. *Eur. Food Res. Technol.*, 2002, **215**, 380–383.
43. Stratton, J.E., Hutkins, W.R. and Taylor, S.L., Biogenic amines in cheese and other fermented foods – a review. *J. Food Protect.*, 1991, **54**, 460–470.
44. Tailor, S. A.N., Shulman, K.I., Walker, S.E., Moss, J. and Gardner, D., Hypertensive episode associated with phenelzine and tap beer – a reanalysis of the role of pressor amines in beer. *J. Clin. Psychopharmacol.*, 1994, **14**, 5–14.
45. Teti, D., Visalli, M. and McNair, H., Analysis of polyamines as markers of (patho)physiological conditions. *J. Chromatogr. B*, 2002, **781**, 107–149.
46. Vally, H. and Thompson, P.J., Allergic and asthmatic reactions to alcoholic drinks. *Addict. Biol.*, 2003, **8**, 3–11.
47. Vidaud, Z., González, E. and García Roché, M.O., Some considerations about the tyramine content of some Cuban beers and wines. *Nahrung*, 1989, **33**, 793–794.
48. Zhang, L.Y., Huang, W.H., Wang, Z.L. and Cheng, J., Determination of histamine by capillary zone electrophoresis with amperometric detection. *Anal. Sci.*, 2002, **18**, 1117–1120.
49. Zimmermeier, S., Vogt, M., Kunz, B., Investigations about amine formation by different microorganisms isolated from wheat-beer. *Monatsschr. Brauwiss.*, 1998, **51**, 82–88 (in German).

(Manuscript accepted for publication May 2003)